

Rooted Clusters for Graph LP Algebras

Esther Banaian^{*1}, Sunita Chepuri^{†2}, Elizabeth Kelley^{‡3}, and
Sylvester W. Zhang^{§1}

¹ Department of Mathematics, University of Minnesota, Minneapolis, MN, USA

² Department of Mathematics, University of Michigan, Ann Arbor, MI, USA

³ Department of Mathematics, University of Illinois Urbana-Champaign, Champaign, IL, USA

Abstract. LP algebras, introduced by Lam and Pylyavskyy, are a generalization of cluster algebras. These algebras are known to have the Laurent phenomenon, but positivity remains conjectural. Graph LP algebras are finite LP algebras encoded by a graph. For the graph LP algebra defined by a tree, we define a family of clusters called *rooted clusters*. We prove positivity for these clusters by giving explicit formulas for each cluster variable. We also give a combinatorial interpretation for these expansions using a generalization of T -paths.

Keywords: Cluster Algebras, Laurent Phenomenon Algebras, Positivity

1 Introduction

Cluster algebras were introduced by Fomin and Zelevinsky [2] as a tool for studying total positivity and dual canonical basis of algebraic groups, and have since been linked to quiver representations, integrable systems, Poisson geometry, Teichmüller theory, mathematical physics, and other topics. They are commutative rings with a family of distinguished generators called *cluster variables*. The cluster variables occur in overlapping subsets of fixed size called *clusters*. Given a cluster \mathcal{C} , we can obtain a unique distinct cluster \mathcal{C}' by a process called *mutation* where one cluster variable in \mathcal{C} is replaced with a different cluster variable. The two cluster variables involved in this process are related by a *binomial exchange relation*; that is, their product can be expressed as a binomial in terms of the other variables in \mathcal{C} (or, equivalently, in \mathcal{C}').

Cluster algebras have several important features, including the following.

- (1) (*Laurent phenomenon*) Given a fixed choice of cluster $\mathcal{C} = (x_1, \dots, x_n)$, every cluster variable can be written as a Laurent polynomial in x_1, \dots, x_n .

*banai003@umn.edu.

†chepu003@umn.edu. Sunita Chepuri was partially supported by NSF Grant No. DMS-1439786.

‡kelleye@illinois.edu. Elizabeth Kelley was partially supported by NSF Grant No. DMS-1937241.

§swzhang@umn.edu.

(2) (*Positivity*) The Laurent polynomial in (1) has positive coefficients.

Lam and Pylyavskyy introduced Laurent phenomenon (LP) algebras as a generalization of cluster algebras [3]. In an LP algebra, the restriction that exchange relations be binomial is relaxed to allow arbitrary irreducible polynomials. Lam and Pylyavskyy proved that the Laurent phenomenon holds for LP algebras and conjectured that positivity holds as well.

Graph LP algebras are a particularly nice class of LP algebras having exchange relations that can be encoded in a graph. Lam and Pylyavskyy explored graph LP algebras in depth in [4] and gave simple descriptions of all of the clusters along with several formulas for computing the cluster variables. However, positivity for graph LP algebras remains conjectural. In this paper, we describe progress towards that conjecture.

We begin by introducing *rooted clusters* for graph LP algebras. Our first main result, which we prove by giving explicit formulas for every cluster variable in terms of each rooted cluster, is positivity for such clusters.

Theorem 1. *If Γ is a tree and \mathcal{C} is a rooted cluster for Γ , then every cluster variable in the graph LP algebra associated to Γ can be expressed as a Laurent polynomial with positive coefficients in the elements of \mathcal{C} .*

We then introduce a generalization of Schiffler’s T -paths for type A cluster algebras [7] for our setting.

Theorem 2. *Let Γ be a tree and \mathcal{C} be a rooted cluster for Γ . If S is a connected subset of vertices of Γ , then the cluster variable Y_S has the combinatorial expansion formula*

$$Y_S = \sum_{\substack{\text{complete hyper} \\ T\text{-paths } \alpha \text{ for } S}} \text{wt}(\alpha).$$

We will begin in [Section 2](#) by giving more background graph on LP algebras and introducing rooted clusters. [Section 3](#) gives formulas for the cluster variables in terms of a rooted cluster \mathcal{C} and sketches the proof of [Theorem 1](#). We begin [Section 4](#) with background on T -paths for type A cluster algebras and then define hyper T -paths. This section culminates with a summary of the proof of [Theorem 2](#). We conclude with a few thoughts about future work.

2 Preliminaries

Laurent phenomenon (LP) algebras were defined by Lam and Pylyavskyy in [3]. A seed for an LP algebra consists of n cluster variables each with an associated exchange polynomial. We can *mutate* at any cluster variable to obtain a new seed. An LP algebra

is a ring generated by all the cluster variables we can obtain from some initial seed by mutation. This paper will focus on a subset of LP algebras called *graph LP algebras*. As these LP algebras can be defined in an equivalent and simpler way (see [Theorem 3](#)), we will not include a full definition of LP algebras here and instead direct interested readers to [3].

For every undirected graph Γ , we obtain a graph LP algebra \mathcal{A}_Γ . The initial seed for \mathcal{A}_Γ is encoded by the edges of the graph.

Definition 1. Let Γ be an undirected graph on $[n] := \{1, \dots, n\}$ and $R = \mathbb{Z}[A_1, \dots, A_n]$. Then the *graph LP algebra* \mathcal{A}_Γ is the LP algebra generated by initial seed

$$\left\{ \left(X_i, A_i + \sum_{i \text{ adjacent to } j} X_j \right) \right\}_{1 \leq i \leq n}$$

Lam and Pylyavskyy prove that these LP algebras have a particularly nice structure using nested collections.


Definition 2. Let Γ be an undirected graph on $[n]$. A family of subsets of $[n]$, $\mathcal{S} = \{S_1, \dots, S_k\}$, is a *nested collection* if

- for any $i, j \leq k$, either $S_i \subseteq S_j$, $S_j \subseteq S_i$, or $S_i \cap S_j = \emptyset$, and
- if $S_{i_1}, \dots, S_{i_\ell}$ are pairwise disjoint, then $S_{i_1}, \dots, S_{i_\ell}$ are exactly the connected components of $\bigcup_{j=1}^\ell S_{i_j}$.

We say \mathcal{S} is a *maximal nested collection* on S if $\bigcup_{i=1}^k S_i = S$ and there is no $S' \subseteq S$ such that $\{S_1, \dots, S_k, S'\}$ is a nested collection.

If Γ is a graph on $[n]$ and \mathcal{S} is a maximal nested collection on $S = [n]$, we will generally say that \mathcal{S} is a maximal nested collection without specifying S .

Example 1. Let Γ be the following graph:



Then $\mathcal{S} = \{\{1\}, \{3\}, \{1, 2, 3, 4\}\}$ is a nested collection on $S = \{1, 2, 3, 4\}$. However, it is not maximal because adding the set $S' = \{1, 2, 3\}$ still yields a nested collection.

As a nonexample, consider $\mathcal{S} = \{\{1\}, \{1, 2\}, \{3\}, \{1, 2, 3, 4\}\}$. We can see that this is not a nested collection by looking at the disjoint sets $\{1, 2\}$ and $\{3\}$. The union of these sets is $\{1, 2, 3\}$, which has only one connected component.

Theorem 3 (Theorem 1.1 of [4]). Let Γ be an undirected graph on $[n]$. Define the matrix $\mathfrak{N} = (n_{ij})$ by

$$n_{ij} = \begin{cases} \frac{A_i + \sum_{i \text{ adjacent to } j} X_j}{X_i} & i = j, \\ -1 & i \text{ adjacent to } j, \\ 0 & \text{otherwise.} \end{cases}$$

Then the graph LP algebra \mathcal{A}_Γ has cluster variables $\{X_1, \dots, X_n\} \cup \{Y_S \mid S \subset [n] \text{ is connected}\}$ where Y_S is the determinant of the submatrix of \mathfrak{N} obtained by taking only rows and columns indexed by S . The clusters for \mathcal{A} are of the form $\{X_{i_1}, \dots, X_{i_k}\} \cup \{Y_S \mid S \in \mathcal{S}\}$ where \mathcal{S} is a maximal nested collection on $[n] \setminus \{i_1, \dots, i_k\}$.

In a slight abuse of notation, we will generally write $Y_{s_1 \dots s_r}$ as shorthand for $Y_{\{s_1, \dots, s_r\}}$.

Example 2. Let Γ be the graph from [Example 1](#). One example of a valid cluster for \mathcal{A}_Γ is $\{X_2, Y_1, Y_5, Y_{35}, Y_{345}\}$. This is because $\{\{1\}, \{5\}, \{3, 5\}, \{3, 4, 5\}\}$ is a maximal nested collection on $\{1, 3, 4, 5\}$. In this case, the \mathfrak{N} matrix is:

$$\mathfrak{N} = \begin{bmatrix} \frac{A_1+X_2}{X_1} & -1 & 0 & 0 & 0 \\ -1 & \frac{A_2+X_1+X_3}{X_2} & -1 & 0 & 0 \\ 0 & -1 & \frac{A_3+X_2+X_4+X_5}{X_3} & -1 & -1 \\ 0 & 0 & -1 & \frac{A_4+X_3}{X_4} & 0 \\ 0 & 0 & -1 & 0 & \frac{A_5+X_3}{X_5} \end{bmatrix}$$

We can use this to rewrite the Y -variables in our cluster. For example,

$$Y_{35} = \begin{vmatrix} \frac{A_3+X_2+X_4+X_5}{X_3} & -1 \\ -1 & \frac{A_5+X_3}{X_5} \end{vmatrix} = \frac{A_3A_5 + A_5X_2 + A_5X_4 + A_5X_5 + A_3X_3 + X_2X_3 + X_3X_4}{X_3X_5}$$

Lam and Pylyavskyy also completely describe the exchange relations for \mathcal{A}_Γ (see Lemmas 4.7 and 4.11 of [4]).

We focus on the case when Γ is a tree. In this setting, we can define a special type of cluster we call a *rooted cluster* that has desirable properties. There is one rooted cluster \mathcal{C}_v for each vertex v of Γ . In order to define this cluster, we think of Γ as being rooted at v . We then think of Γ as a poset with the root v being the maximal element and cover relations given by edges of Γ . This leads us to establish the following notation:

- Notice that if $i \neq v$, then i is covered by exactly one vertex. We call this vertex i^+ .
- The set of elements covered by i is denoted $\Gamma_{<i}^v$. Similarly we have the sets $\Gamma_{>i}^v$, $\Gamma_{<i'}^v$, $\Gamma_{>i'}^v$, and $\Gamma_{\geq i}^v$ (note that $\Gamma_{>i}^v = \{i^+\}$ if $i \neq v$).




Figure 1: We can picture the cluster rooted at vertex 3 by taking subsets which are closed going down. For example, $I_2 = \{1, 2, 5\}$ and $I_5 = \{5\}$.

Definition 3. Let Γ be a tree on $[n]$. Make Γ into a rooted tree by choosing a vertex v to be the root. Then for each vertex x in Γ , let $I_x = \Gamma_{\leq x}^v$. The *rooted cluster* \mathcal{C}_v is $\{I_x\}_{x \in [n]}$.

Example 3. Consider the tree in **Figure 1**. The cluster \mathcal{C}_3 consists of $I_1 = \{1\}$, $I_2 = \{1, 2, 5\}$, $I_3 = \{1, 2, 3, 4, 5\}$, $I_4 = \{4\}$, and $I_5 = \{5\}$. One can check that these sets form a nested collection, and that it is not possible to add another set compatible with all others. If we root at 2 instead, then \mathcal{C}_2 consists of the sets $I_1 = \{1\}$, $I_2 = \{1, 2, 3, 4, 5\}$, $I_3 = \{3, 4\}$, $I_4 = \{4\}$, and $I_5 = \{5\}$.

3 Formulas

Although **Theorem 3** and the exchange relations for graph LP algebras give formulas for all cluster variables, these formulas are not guaranteed to be in terms of the variables of any particular cluster. In this section, we provide formulas for each of the X_i and Y_S cluster variables in terms of a rooted cluster \mathcal{C}_v . Positivity for this case follows as an immediate consequence.

One useful observation, which simplifies the presentation of our formulas, is that:

Lemma 1. Let Γ be a tree rooted at v and i a vertex of Γ . Then

$$Y_{\Gamma_{<i}^v} = \prod_{u \in \Gamma_{<i}^v} Y_{I_u}$$

and for $u \in \Gamma_{<i}^v$ we have,

$$Y_{\Gamma_{<i}^v \setminus \Gamma_{\geq u}^v} = \prod_{w \in \Gamma_{<i}^v \cap \Gamma_{\geq u}^v} \prod_{x \in \Gamma_{<w}^v \setminus \Gamma_{\geq u}^v} Y_{I_x}$$

Using **Theorem 3** and Lemmas 4.7 and 4.11 of [4], we obtain an expansion formula for Y_S when $|S| = 1$. It follows from **Lemma 1** that this formula only includes Y -variables from \mathcal{C}_v .

Proposition 1. Let Γ be a tree rooted at v . For every vertex i of Γ ,

$$Y_{\{i\}} = \frac{Y_{I_i} + \sum_{u \in \Gamma_{<i}^v} Y_{\Gamma_{<i}^v \setminus \{u\}}}{Y_{\Gamma_{<i}^v}}.$$

Example 4. Consider the tree shown in [Figure 1](#) and the rooted cluster \mathcal{C}_3 . Using the formula from Proposition 1, we compute

$$Y_{\{2\}} = \frac{Y_{I_2} + \sum_{u \in \Gamma_{<2}^3} Y_{\Gamma_{<2}^3 \setminus \{u\}}}{Y_{\Gamma_{<2}^3}} = \frac{Y_{I_2} + Y_{\Gamma_{<2}^3 \setminus \{1\}} + Y_{\Gamma_{<2}^3 \setminus \{5\}}}{Y_{\Gamma_{<2}^3}} = \frac{Y_{\{1,2,5\}} + Y_5 + Y_1}{Y_1 Y_5}.$$

Later, we will see that this is consistent with the expansion computed via our hyper T -path construction.

We then establish the following technical lemmas, which are used to obtain expansion formulas for Y_S for arbitrary S .

Lemma 2. Let Γ be a tree, S be a connected subset of vertices, and $T = \{(a, b) \in S \times S \mid a = b^+\}$. Then

$$Y_S = \sum_{n=0}^{\lfloor \frac{|S|}{2} \rfloor} (-1)^n \sum_{A \in A(n)} \left(\prod_{x \in (S \setminus A')} Y_{\{x\}} \right)$$

where $A(n) = \{A \subseteq 2^T \mid |A| = n \text{ and } \{a, b\} \cap \{c, d\} = \emptyset \text{ for all } (a, b), (c, d) \in A \text{ with } (a, b) \neq (c, d)\}$ and for any $A \in A(n)$, $A' = \{s \mid s \text{ is part of a pair in } A\}$.

Lemma 3. Let $Y_S^{(n)} = \sum_{A \in A(n)} \left(\prod_{x \in (S \setminus A')} Y_{\{x\}} \right)$. If t is a monomial that appears as a term in $Y_S^{(m+1)}$, then t also appears in $Y_S^{(m)}$.

As a consequence of [Lemma 3](#), the set of monomials that appear in the expansion of the sum in [Lemma 2](#) are exactly those that appear in the expansion of $\prod_{x \in S} Y_{\{x\}}$. By counting the number of times each monomial appears in the expansion of the sum in [Lemma 2](#), we obtain an expansion formula for all Y_S . To obtain an expansion formula for all X_i , we first apply Lemmas 4.7 and 4.11 of [\[4\]](#) to $S = \Gamma_{<i}^v$ to obtain a formula for X_v and then induct on the distance from i to the root vertex v .

Theorem 4. Let Γ be a tree rooted at v . For all vertices i of Γ and vertex subsets S , we have

$$X_i = \frac{\sum_{u \in \Gamma_{\geq i}^v} \left(\prod_{w \in \Gamma_{\geq i}^v \setminus \Gamma_{\geq u}^v} Y_{\Gamma_{<w}^v} \right) \left(\prod_{w \in \Gamma_{> u}^v} Y_{I_w} \right) \left(\sum_{w \in I_u} Y_{\Gamma_{<u}^v \setminus \Gamma_{\geq w}^v} A_w \right)}{\prod_{u \in \Gamma_{\geq i}^v} Y_{I_u}}, \quad (3.1)$$

$$Y_S = \sum_{\substack{O \subseteq S \text{ containing all} \\ \text{minimal elements of } \Gamma \text{ in } S}} \sum_{\substack{u: S \setminus O \rightarrow V(\Gamma) \\ u(x) \in \Gamma_{<x}^v \setminus O}} \frac{\left(\prod_{x \in O} Y_{I_x} \right) \left(\prod_{x \in S \setminus O} Y_{\Gamma_{<x}^v \setminus \{u(x)\}} \right)}{\prod_{x \in S} Y_{\Gamma_{<x}^v}}. \quad (3.2)$$

[Theorem 1](#) follows as a corollary of these expansion formulas, using [Lemma 1](#).

4 Hyper T -paths

We reprove [Theorem 1](#) about positivity of the Y -variables via a combinatorial interpretation of the expansions of Y_S in terms of a rooted cluster. Our constructions are a generalization of T -paths, which were originally introduced by Schiffler [7] as a tool for finding cluster expansion formulas in type A cluster algebras.

4.1 T -paths for Type A Cluster Algebras

Type A cluster algebras are modeled by triangulations of an $(n + 3)$ -gon, with each initial seed corresponding to a unique initial triangulation. Consider an $(n + 3)$ -gon with vertices labeled $1, \dots, n + 3$ and a fixed triangulation $T = \{T_1, \dots, T_n, T_{n+1}, \dots, T_{2n+3}\}$ where T_1, \dots, T_n are interior diagonals and T_{n+1}, \dots, T_{2n+3} are boundary edges. Let i and j be non-adjacent boundary vertices and let $M_{i,j}$ denote the interior diagonal connecting i and j . Fix an orientation on $M_{i,j}$ and let $i = p_0, p_1, \dots, p_d, p_{d+1} = j$ be the ordered list of intersection points of $M_{i,j}$ and arcs of T . Then let i_1, \dots, i_d be a list of indices such that intersection point p_k lies on the arc $T_{i_k} \in T$. For $k \in [d]$, let M_k denote the segment of the diagonal $M_{i,j}$ between the intersection points p_k and p_{k+1} .

In [5], Musiker and Schiffler define a *complete T -path from i to j* as a sequence $\alpha = (t_1, \dots, t_{\ell(\alpha)})$ such that

- (T1) $i = a_0, a_1, \dots, a_{\ell(\alpha)} = j$ are (not necessarily distinct) vertices of the $(n + 3)$ -gon,
- (T2) $t_k \in \alpha$ is an arc in the triangulation T that connects vertices a_{k-1} and a_k , and
- (T3) the even arcs are precisely the arcs crossed by $M_{i,j}$ in order, i.e. $t_{2k} = T_{i_k}$.

We denote the set of all complete T -paths from i to j as \mathcal{T}_{ij} . Given a complete T -path α , the *weight* of α is defined to be the Laurent monomial

$$\text{wt}(\alpha) := \left(\prod_{i \text{ odd}} \text{wt}(t_i) \right) \left(\prod_{i \text{ even}} \text{wt}(t_i) \right)^{-1}$$

where the weight of edge t_i is given by $\text{wt}(t_i) := x_{t_i}$. By summing over the set \mathcal{T}_{ij} , Schiffler [7] then obtains an expansion formula for the cluster variable corresponding to $M_{i,j}$ in terms of the cluster seed corresponding to the triangulation T :

$$x_{M_{i,j}} := \sum_{\alpha \in \mathcal{T}_{ij}} \text{wt}(\alpha)$$

Although it is not immediately obvious, this cluster expansion formula is independent of the choice of orientation on $M_{i,j}$.

4.2 Construction and Examples

In this section, we generalize the notion of T -paths to define *hyper T -paths*. We first construct an auxiliary graph, $\Gamma_{\mathcal{C}}$.

Let Γ be a tree and \mathcal{C} be a rooted cluster for Γ . For each vertex x of degree 1 in $\Gamma_{\mathcal{C}}$ we add an additional vertex x' which is adjacent only to x . Call this extended graph Γ' . We will continue to think of Γ' as a poset where $x' < x$ if x is not the root and $x' > x$ if x is the root. For every $S \in \mathcal{C}$, let S' be the set of vertices in Γ' that are adjacent to a vertex in S but are themselves not in S . Add a hyperedge labelled S which connects all the vertices of S' . As a convenient abuse of notation, we often refer to this hyperedge simply as S . We refer to this new hypergraph as $\Gamma_{\mathcal{C}}$. See [Figure 2](#) for an example.




Figure 2: The hypergraph associated to the tree from [Figure 1](#).

For x a vertex in the rooted tree Γ , we will use \mathcal{L}_x to denote to all minimal elements of Γ' that are less than x . Equivalently, \mathcal{L}_x is the elements of $\Gamma' \setminus \Gamma$ that are less than x .

Definition 4. Let S be a connected subset of Γ . A *complete hyper T -path* for S with respect to \mathcal{C} is a set of nodes, labelled by vertices of $\Gamma_{\mathcal{C}}$, joined by connections labelled by hyperedges of $\Gamma_{\mathcal{C}}$ such that the diagram is connected and the following hold.

1. If a connection is labelled by hyperedge e , it joins nodes labelled by all the endpoints of e with multiplicity 1.
2. There are a distinguished set of *boundary nodes* labelled by elements of S' with multiplicity 1. Other nodes are called *internal nodes*.
3. Connections are specified to be *even* or *odd*.
4. Boundary nodes are adjacent only to odd connections.
5. Internal nodes labelled by elements of S are adjacent to exactly one even and at least one odd connection.


6. Internal nodes labelled by elements not in S are adjacent to exactly one even and exactly one odd connection.
7. If x, y are below elements of S , any path in any complete hyper T -path from boundary node x to boundary node y uses even connections labelled, in order, by $I_x, I_{a_p}, I_{a_{p-1}}, \dots, I_{a_1}, I_{b_1}, I_{b_2}, \dots, I_{b_q}, I_y$ where the shortest path from x to y in Γ' is $x, a_p, a_{p-1}, \dots, a_1, x \vee y, b_1, b_2, \dots, b_q, y$ for $p, q \geq 0$.
8. If x is below an element of S and y above the maximal element of S , any path in any complete hyper T -path from the boundary node x to the boundary node y uses even connections labelled, in order, by $I_x, I_{a_p}, \dots, I_{a_2}$, where the shortest path from x to y in Γ' is $x, a_p, a_{p-1}, \dots, a_1, y$, $p \geq 1$. If $p = 1$, then a path from x to y uses the even connection I_x .
9. If x, y are boundary nodes, where the shortest path from x to y in Γ' is $x, a_p, \dots, a_1, x \vee y, b_1, \dots, b_q, y$, then any path in any complete hyper T -path from x to y uses nodes labelled by elements of $\mathcal{L}_{x \vee y}$ and $a_p, a_{p-1}, \dots, a_1, x \vee y, b_1, b_2, \dots, b_q$, with any multiplicity. If one of the nodes, say y , is adjacent to the maximal element of S , then $x \vee y = y$ and $q = 0$.

Definition 5. The *weight* of a hyper T -path α is

$$\text{wt}(\alpha) = \left(\prod_{\text{odd connections } c} \text{wt}(c) \right) \left(\prod_{\text{even connections } c} \text{wt}(c) \right)^{-1}$$


where the weight of a connection labelled by a set I_x is Y_{I_x} and the weight of a connection labelled by an edge in Γ' is 1.

Example 5. Consider the graph Γ in [Figure 1](#) with a cluster rooted at 3. Below we list the three hyper T -paths associated to the set $\{2\}$.



From left to right, these T -paths have weight $\frac{1}{Y_{I_5}}, \frac{1}{Y_{I_1}}$ and $\frac{Y_{I_2}}{Y_{I_1}Y_{I_5}}$. This is consistent with the expansion of Y_2 computed by [Proposition 1](#), shown in [Example 4](#).

We also give one example of a T -path associated to the set $\{2, 3\}$ from the same graph with the same rooted cluster. The T -path below has weight $\frac{1}{Y_{I_1}Y_{I_4}}$.



Remark 1. The T -paths in [Example 5](#) illustrate that these sometimes have adjacent odd and even connections with the same label. The contributions of these connections then cancels in the weight of the T -path. To avoid this, one could instead define *reduced hyper T -paths*. Here, we find it simpler to work with complete hyper T -paths.

Our main result in this section ([Theorem 2](#)) is that the hyper T -paths provide a combinatorial interpretation of the expansion of a variable Y_S with respect to a rooted cluster. Because the coefficients of this expansion formula count the number of hyper T -paths with a particular weight, they are manifestly positive integers. By [Equation \(3.1\)](#), this also implies that the coefficients of the X -variables are positive; thus, the hyper T -paths provide a second proof of [Theorem 1](#).

The proof of [Theorem 2](#) has similarities with the proof of [Theorem 4](#). We first prove the theorem in the case of singleton sets; recall the description of the expansion of $Y_{\{i\}}$ in [Proposition 1](#). We then define valid *pastings* of hyper T -paths and study which T -paths associated to $\{i\}$ and $\{j\}$ can be pasted, provided that i and j are adjacent in Γ . The pasting requirements mirrors the algebraic computations from the proof of [Theorem 2](#), leading us to our result. See [\[1\]](#) for complete proofs of both results.

5 Future Directions

5.1 From Rooted Clusters to Other Clusters

We would like to extend our results to other clusters for trees. Our definition of rooted clusters is motivated by the algebraic formulas for the exchange relations (Lemmas 4.7 and 4.11 of [\[4\]](#)). For rooted clusters, the exchange relations give expansions in terms of the cluster variables in the initial seed. This is not, however, true for an arbitrary cluster.

Proving formulas algebraically for other types of clusters will likely require an inductive argument, which seems to be easiest in star graphs because of their symmetry.

Conjecture 1. Let S_n denote the star graph on n vertices whose central vertex is labeled by 1. Let \mathcal{C} be the cluster $\{\{3\}, \{4\}, \dots, \{n\}, \{1, 3, 4, \dots, n\}\}$. For any vertex subset S such that $1 \in S$, we conjecture that

$$Y_S = \begin{cases} \frac{\prod_{i \in S \setminus \{1,2\}} Y_i^2}{Y_{[n] \setminus \{1,2\}}} \left(\sum_{i \notin S} \prod_{j \neq 1,2,i} Y_j \right) + \frac{Y_{[n]}}{\prod_{i \in [n] \setminus S} Y_i} + \frac{Y_{[n]}}{Y_{[n] \setminus \{2\}}} \left(\sum_{i \in [n] \setminus S} \frac{1}{Y_i} \right) & \text{if } 2 \notin S \\ \frac{Y_{[n] \setminus \{2\}}}{\prod_{i \in [n] \setminus (S \cup \{2\})} Y_i} + \left(\prod_{i \in S \setminus \{1\}} Y_i \right) \left(\sum_{j \in [n] \setminus (S \cup \{2\})} \frac{1}{Y_j} \right) & \text{if } 2 \in S \end{cases}$$

If $1 \notin S$, then S either consists of a set of disconnected leaves or a single leaf. Because $\{i\} \in \mathcal{C}$ for all $i \neq 2$, we then have

$$Y_s = \prod_{i \in S} Y_i$$

We are also hopeful that we can extend our hyper T -path expansion formula to other types of clusters. For a type A cluster algebra, T -paths can be used to find expansions for cluster variables in terms of any cluster. Because the definitions of T -paths and hyper T -paths for path graphs are similar, this suggests that it might be possible to use hyper T -paths for other clusters when Γ is an arbitrary tree.

Unfortunately, our current hyper T -path construction does not work for arbitrary clusters. One immediate problem is that Rules (7) and (8) would need to be rewritten to be more general. That is, we would need to allow the even edges to be labelled by any set in the cluster that is incompatible with S . However, that change would be insufficient because we still wouldn't have "enough" valid hyper T -paths. Further, the expansions of some cluster variables with respect to certain clusters contain monomials with squared terms in the denominator. This is not possible with our current definition. Thus, it is clear that there is some other modifications are required for us to be able to extend our construction to other clusters.

5.2 Snake Graphs

In [6], Musiker, Schiffler and Williams provided an alternative combinatorial formula for type A cluster algebras using perfect matchings on certain *snake graphs*. For such cluster algebras, there exists a weight-preserving bijection between the set of (complete) T -paths associated to an arc and the set of perfect matching of its snake graph. Since our hyper T -path are generalizations of complete T -paths, we hope to extend this bijection to obtain a graph-theoretic formula analogous to [6] for LP algebras from trees. We illustrate the rough idea in the following example.




Figure 3: A generalized snake graph and one of its matchings.

Example 6. On the left of Figure 3, we draw a snake graph associated to the set $\{2, 3\}$ of the graph Γ in Figure 2. In its perfect matchings, the vertex 2 is allowed to have valence 2. On the right of Figure 3 is the perfect matching which has the same weight as the hyper T -path in Example 5.

Snake graphs may prove more tractable for non-rooted clusters; however, the full construction of a generalized snake graph is still in progress. We were able to complete the preceding example because all vertices in the underlying graph Γ had degree three or less. When the vertices in Γ have higher degree, it is unclear how to draw the snake graphs.

Acknowledgements

We would like to thank Pavlo Pylyavskyy for suggesting this problem, Trevor Karn for organizing the Minnesota Combinatorics Working Group, and Kayla Wright and Libby Farrell for generating initial computational data.

References

- [1] Esther Banaian et al. “Rooted clusters for graph LP algebras”. In: *arXiv preprint arXiv:2107.14785* (2021).
- [2] Sergey Fomin and Andrei Zelevinsky. “Cluster algebras I: foundations”. In: *Journal of the American Mathematical Society* 15.2 (2002), pp. 497–529.
- [3] Thomas Lam and Pavlo Pylyavskyy. “Laurent phenomenon algebras”. In: *arXiv preprint arXiv:1206.2611* (2012).
- [4] Thomas Lam and Pavlo Pylyavskyy. “Linear Laurent phenomenon algebras”. In: *International Mathematics Research Notices* 2016.10 (2016), pp. 3163–3203.
- [5] Gregg Musiker and Ralf Schiffler. “Cluster expansion formulas and perfect matchings”. In: *Journal of Algebraic Combinatorics* (2009).
- [6] Gregg Musiker, Ralf Schiffler, and Lauren Williams. “Positivity for cluster algebras from surfaces”. In: *Advances in Mathematics* 227.6 (2011), pp. 2241–2308.
- [7] Ralf Schiffler. “A cluster expansion formula (A_n case)”. In: *The Electronic Journal of Combinatorics* 15.R64 (2008).